



Occupational &
Environmental Medicine

Musculoskeletal Conditions and Chronic Pain Among Working Patients

Andrea Furlan

Scientist

Institute for Work & Health

Lynn Cooper

Director of Research & Education
Canadian Injured Workers Alliance

November 3, 2023

Faculty/Presenter Disclosure

- **Faculty:** Andrea Furlan, MD PhD
- **Relationships with financial sponsors:**
 - **Grants/Research Support:** Health Canada, CIHR, Ontario Health, Canadian Generic Products Association, WSIB Grants Program, WorkSafe BC, Desjardins Insurance
 - **Speakers Bureau/Honoraria:** No
 - **Consulting Fees:** WSIB Drug Advisory Committee and Health Advisory Committee
 - **Patents:** No
 - **Other:** YouTube monetized channel, Amazon affiliate marketing, Book Royalties
Scientist, Institute for Work & Health; Psychiatrist, Senior Scientist, Toronto Rehabilitation Institute, UHN; Associate Professor, Division of Psychiatry, Department of Medicine, University of Toronto; Co-chair ECHO Ontario Occupational and Environmental Medicine

Faculty/Presenter Disclosure

- **Presenter:** Lynn Cooper, BES
Lived Experience – work injury & persistent pain
- **Relationships with financial sponsors:**
 - **Grants/Research Support:** No
 - **Speakers Bureau/Honoraria:** Dalhousie, Manitoba, McMaster
 - **Consulting Fees:** No
 - **Patents:** No
 - **Other:** Employee of Canadian Injured Workers Alliance

Disclosure of Financial Support

- This program has received financial support from the Workplace Safety and Insurance Board (WSIB) in the form of an educational grant.
- This program has received in-kind support from – N/A
- Potential for conflict(s) of interest:
 - None



Mitigating Potential Bias



Occupational &
Environmental Medicine

- The information presented in this program is based on recent information that is explicitly “evidence-based”.
- This Program and its material is peer reviewed and all the recommendations involving clinical medicine are based on evidence that is accepted within the profession; and all scientific research referred to, reported, or used in this CME/CPD activity in support or justification of patient care recommendations conforms to the generally accepted standards

Learning Objectives

By the end of this session, participants will be able to:

1. Describe an approach to examine a person with a musculoskeletal (MSK) problem
2. Explain the indications of opioids for nociceptive, neuropathic and nociplastic chronic pain
3. Cite 10 evidence-based treatments for low back pain
4. Understand the impact of working with chronic pain on the person






Fig. 2. Stakeholders' rating of factors influencing return to work. RTW = return to work; UE = upper extremity; HCP = health-care provider; SDP = suitable duties program; WC = workers' compensation.

Prevalence of Disability by Age Group, Canada

Prevalence of Disabilities by age group, Canada 2012



Prevalence of Disabilities by age group, Canada 2012



The most prevalent underlying pain-related conditions reported by those with pain-related disabilities were arthritis, dorsalgia, and dorsopathy.

MSK Lesions

Strain	
Sprain	
Contusion	

Dislocation	
Subluxation	

Synovitis	
Bursitis	

Rupture	
Tear	

Tendinopathy	
Tenosynovitis	
Tendinitis	
Calcific tendinitis	
Tendinosis	

Overuse syndrome	
Cumulative trauma disorder	
Repetitive strain injury	

MSK Lesions

Strain	Overexertion in a muscle/tendon Grades: I (mild), II or III (rupture)
Sprain	Injury to a ligament Grades: I (mild), II or III (rupture)
Contusion	Capillary rupture, bleeding

Dislocation	Displacement with soft tissue damage
Subluxation	Partial dislocation

Synovitis	Inflammation of synovial membrane
Bursitis	Inflammation of a bursae

Rupture	Rupture and Tear are synonyms. Partial = pain; Complete = painless
Tear	

Tendinopathy	General term for tendon injury
Tenosynovitis	Inflammation of synovial membrane covering a tendon
Tendinitis	Inflammation of tendon
Calcific tendinitis	Tendinitis with calcium deposit
Tendinosis	Degeneration due to repetitive microtrauma

Overuse syndrome	Repeated, <u>submaximal</u> overload and/or frictional wear to a muscle or tendon resulting in inflammation and pain.
Cumulative trauma disorder	
Repetitive strain injury	


MSK Lesions – Time to Heal

	weeks						months						
	1	2	3	4	5	6	2	4	6	8	10	12	18
Muscle Tendon Ligament injury	Grade 1 (mild)						Grade II		Grade III (tear)				

Shoulder	Subluxation													
							Dislocation							
									Frozen Shoulder					
	Forming calcific tendinitis (no pain)						Little pain		Severe pain		No pain			

Nerve				Post carpal tunnel release									
				Sciatica									

Tousignant-Laflamme, 2017
J Pain Research



Nociplastic pain is defined as 'Pain that arises from altered nociception despite no clear evidence of actual or threatened tissue damage causing the activation of peripheral nociceptors or evidence for disease or lesion of the somatosensory system causing the pain (IASP)




3 types of pain

- Nociceptive
 - Neuropathic
 - Nociplastic
- Kosek, 2016 Do we need a third mechanistic description for chronic pain states?


What is Pain?

Pain is "an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage." (IASP 2020)

Descartes, 1644



What is Nociceptive Pain?



Chronic Pain is a Disease


Chronic pain was recently recognized by the World Health Organization (WHO) as a disease in its own right, resulting in revisions to the latest (11th) version of the International Classification of Diseases (ICD-11).

According to ICD-11, chronic pain can be further classified as **chronic primary pain** or chronic secondary pain.


Chronic primary pain is pain in one or more anatomical regions that:

1. Persists or recurs for longer than 3 months; and,
2. Is associated with significant emotional distress (e.g., anxiety, anger, frustration, depressed mood) and/or significant functional disability (interference in activities of daily life and participation in social roles); and,
3. The symptoms are not better accounted for by another diagnosis (Nicholas et al., 2019).

Chronic primary pain includes the following sub-diagnoses: chronic widespread pain, complex regional pain syndrome, chronic primary headache or orofacial pain, chronic primary visceral pain, and chronic primary musculoskeletal pain.





Presence of painful symptoms reduce the probability of recovery from depression: 9% versus 47% (Arango-Davila, 2018)



Catastrophizing was the strongest and most consistent psychosocial factor associated with persistence of pain and poor function in persons with chronic pain, even after controlling for depression.

Catastrophizing is a modifiable risk factor (Arango-Davila, 2018)





Flag System for Prognosis in Low Back Pain

	Issue	Description	Actions
Red	Medical Issues	Neurological (cauda equina), Infection, Fracture, Tumour, Inflammation → NIFTI	Admit to hospital Refer to specialist
Orange	Psychiatric Issues	Major personality disorder, Substance Use Disorder, PTSD, Psychosis, High levels of anxiety, distress	Refer to psychiatry consult
Yellow	Psychological Behavioural	Poor coping strategies, Low self-efficacy, Fear avoidance, maladaptive behaviours and beliefs, Family reinforcement, litigation, compensation	Refer to multidisciplinary pain management team
Blue	Perception of Work	Not working, fear of re-injury, poor work satisfaction, work-related stress	Address issues in collaboration with employer
Black	Actual Work Conditions	Poor work conditions, manual work, unsociable hours	Consultation with employer and policy makers

Assessment of Patients with Chronic Pain

Questionnaires

Before the visit

- Screening for depression, anxiety, somatization
- Central Sensitization Inventory
- Neuropathic Pain Screening
- List of current and past treatments (effectiveness and failures)
- Past medical and family history
- Previous investigations/consults

On the day of the visit

- Brief Pain Inventory: pain intensity x 4 and pain interference
- Body pain diagram
- Specific depending on condition (e.g. OHRT if work disability, Fibromyalgia Diagnostic Criteria if widespread pain, or STOP-BANG if sleep apnea).

Clinical Assessment

Patient's goals

History of present illness (SOCRATES*)

Physical examination:

- mental status
- mobility
- focused MSK
- neurological screening

* Site, Onset, Characteristics, Radiating, Associated symptoms, Timing, Exacerbating/Ameliorating, Severity

“5M IS” of Management of Chronic Primary Pain

Mind	Movement	Modalities Manual	Medications	Interventional	Surgery
Pain Reprocessing Therapy (PRT)* Pain Neuroscience Education (PNE)** Graded Motor Imagery (GMI)*** CBT, ACT, MI, Group sessions, Written emotional expression, Psychomotor therapy, MBSR, EMG-biofeedback, distraction, hypnosis,	Aerobics/Cardio Strengthening Stretching Relaxation Water-based Home/group based Pilates Tai Chi Qigong Yoga	Acupuncture Electrotherapy TENS Manipulation Mobilization Massage	Lower dose rational polypharmacy Simple analgesics Serotonin Gaba Tramadol Low-dose naltrexone THC/CBD?	Trigger point injections Nerve blocks Nerve ablation Intra-articular injections Capsular distension Botulinum toxin Regenerative medicine	Joint replacements Spinal cord stimulator Deep brain stimulator Intrathecal pumps

Opioids at the WSIB

REMEMBER: Management of (primary) chronic pain: TAPER OPIOIDS SLOWLY TO THE LOWEST POSSIBLE DOSE (Canadian Opioid Guideline recommendation #9)

Approach based on 2 key principles:


- Authorization of opioids for workers should support treatment goals that include improvement in function, pain relief, quality of life, and safe and sustained return to work
- Management of pain is consistent with current best practice

→ Allow prescriptions for a maximum of 12 weeks

→ Opioid coverage beyond 4 weeks will be subject to clinical review

→ Endorse the 2017 Canadian Opioid Guideline


Thank you



Email afurlan@iwh.on.ca

Twitter [@adfurlan](https://twitter.com/adfurlan)

YouTube [Dr.AndreaFurlan](https://www.youtube.com/DrAndreaFurlan)



PowerOverPain.ca

Extra slides



Occupational &
Environmental Medicine

Chronic Pain and Comorbidities

Insomnia

- Prevalence in the general population:
 - 9% chronic
 - 30% occasional
- Prevalence among persons with chronic pain: 65% to 89%

Anxiety

- Stressful situations in healthy individual → analgesia
- Stressful situations in an individual with central sensitization → hyperalgesia

Not all Chronic Pains are the Same

Without central sensitization (secondary)

- Ascending Pain pathways are intact
- Descending inhibitory pathways are intact
- Underlying chronic pathology - pain
- No signs of central sensitization
- Expected (normal) psychological response
- Its function is to alert the individual to seek treatment
- For example: hip osteoarthritis

“Nociceptive pain”

“Neuropathic pain”

With central sensitization (primary)

- Malfunction of pain system
- No underlying pathology
- Many signs of central sensitization
- Abnormal psychological response to pain
- Difficulty to concentrate, sleep, relationships, work
- Chronic fatigue (physical and mental)
- It has no function to the individual
- For example: fibromyalgia

“Nociplastic pain”

Symptoms of Central Sensitization (CS)



Hypersensitivity to bright light, noise, touch, pesticides, food, mechanical pressure, medication, temperature, weather



Widespread pain



Fatigue (physical and mental)



Sleep disturbance



Numbness



Swelling sensations





Low libido



Low mood

Confirmation with physical exam (sensory examination)



Chronic Pain and Comorbidities

Chronic Pain has worldwide prevalence:
1 in 5 children and adults; 1 in 3 older adults

Pain and Depression

- Prevalence of pain symptoms in patients with depression: 65% (range 15% to 100%)
- Presence of painful symptoms reduce the probability of recovery from depression:
9% versus 47%

Psychology of Pain Catastrophizing

A maladaptive coping style.

A construct with three components:

- magnification or amplification of pain
- ruminating thoughts about pain
- perceived helplessness in the face of pain

The strongest and most consistent psychosocial factor associated with persistence of pain and poor function in persons with chronic pain, even after controlling for depression.

Catastrophizing **is modifiable** and, if treated by psychosocial interventions, pain improves with a decrease in catastrophizing.

Psychology of Pain **Fear Avoidance**

Another maladaptive coping style.

The avoidance of work, movement, or other activities due to fear that they will damage the body or worsen pain.

Pain patients high in fear avoidance have worse long-term outcomes.

Fear avoidance is associated with catastrophic misinterpretations of pain, hypervigilance, increased escape and avoidance behaviors, and increased pain intensity and functional disability.

Pain-related fear may increase the risk for developing new-onset back pain, for its chronification, and for its persistence.

The value of **changing beliefs about pain early in its course** has been shown in studies involving patient education in physician's offices and over the public radio.

Psychology of Pain **Job Satisfaction**

Job satisfaction **is NOT** a prognostic factor for duration of sick leave from chronic pain

Supervisor support **may be a factor** in duration of sick leave from chronic pain

Inconclusive evidence for the effects of job demands, job control, job strain, skill discretion, decision authority, job security, co-worker support, supervisory support, psychological demands, physical demands, and work flexibility on duration of work absenteeism

There is **strong evidence**, however, that heavy work is a predictor for longer duration of sick leave. Although assignment to light duties as commonly used for a rapid return to work appears not to shorten sick leave in workers with acute low back pain, staying active and modified work are supported

Psychology of Pain

Recovery expectation



Occupational &
Environmental Medicine

Strongest predictor of work outcome for patients with pain

Recovery expectations measured within weeks of new-onset of pain can identify people at risk of poor outcome.

Expectation is a complex construct composed of numerous variables such as concerns about pain exacerbations, recurrent pain, financial security, support at work, and self-confidence.

Practitioners may need to further inquire why patients have beliefs of delayed recovery and address specific concerns.